

Curie-Weiss magnet – a simple model of phase transition

Martin Kochmański

University of Rzeszów, Rejtana 16A, 35-310 Rzeszów, Poland

Tadeusz Paszkiewicz and Sławomir Wolski

*Rzeszów University of Technology, ul. Powstańców Warszawy 6, 35-959 Rzeszów, Poland**

(Dated: January 11, 2013)

Abstract

The Curie-Weiss model is an exactly soluble model of ferromagnetism that allows one to study in detail the behavior of thermodynamic quantities in the neighborhood of the critical temperature. Because of its simplicity and because of the correctness of at least of some of its predictions, the Curie-Weiss model occupies an important place in the statistical mechanics literature and its application to information theory. It is frequently presented as an introduction to the Ising model or to spin glass models, and usually only general features of the Curie-Weiss model are presented. We discuss here properties of this model in a rather detailed way. We present the exact, approximate and numerical results for this particular model.

I. INTRODUCTION

A phase transition is the transformation of a thermodynamic system from one phase or state to another. During a phase transition of a given medium certain properties change, often discontinuously, as a result of some external conditions, such as temperature, pressure or the magnetic field. For example, a liquid may become gas upon heating to the boiling point, resulting in an abrupt change in volume.

In the case of a ferromagnet one should predict the dependence of such quantities as the free and internal energy, heat capacity and magnetic susceptibility on temperature T and magnetic induction B . Generally, this is very complicated task. Ma¹ stressed the distinction between the direct approach to the problem of phase transitions and the approach exploiting symmetries of the problem. Here we shall illustrate the former approach. This means calculations of physical properties of interest in terms of parameters given in a particular model, i.e. solving a model. The calculations may be done analytically or numerically; exactly or approximately.

The approach exploiting symmetries does not attempt to solve the model. From various symmetry properties one deduces some important physical properties.

We shall focus on a particular model – the Curie-Weiss model of a magnet. This model is also called the infinite range Ising model. The Curie Weiss model is an exactly soluble model of ferromagnetism that allows one to study in detail the behavior of thermodynamic quantities in the neighborhood of the critical temperature. Since not all predictions of this model agree with experiment, other models must be considered. However, because of its simplicity and because of the correctness of at least some of its predictions, the classical Curie-Weiss model occupies a central place in the statistical mechanics literature. It is frequently presented as an introduction to the Ising model or to spin glass models^{2–10}. In these references only general features of the Curie-Weiss model are discussed. This is why we shall discuss here properties of this model in a rather detailed way. We present the exact, approximate and numerical results for this particular model.

II. THE CURIE-WEISS MODEL

One of simplest classical systems exhibiting phase transition – the Curie-Weiss model has been introduced by Pierre Curie and then by Pierre Weiss in their development of simplified theory of ferromagnetism^{6,9}. We shall call it the C-W magnet. Let us call the set of integers from 1 to N a lattice, and its element i a site. We assign a variable s_i (the Ising spin) to each site. The Ising spin is characterized by the binary value: +1 if microscopic magnetic moment is pointing up or -1 if it is pointed down. Particles with the Ising spins interact via Hamiltonian

$$H_{int} = -\frac{J}{N} \sum_{1 \leq i < j \leq N} s_i s_j. \quad (1)$$

The constant J is positive. The interaction energy of all pairs of spins of the Curie-Weiss magnet is the same and their interaction depends on N . The normalization by $1/N$ makes H_N a quantity of the order N . The underlying assumption of an infinite-range interaction is clearly unphysical. The Hamiltonian (1) does not depend on dimension of the space which Curie-Weiss magnet is occupying.

The magnetic moment of a particle is proportional to the spin $\mu_i = \mu s_i$, where μ is the magnetic moment. In an applied magnetic field with the magnetic induction vector $\mathbf{B}$ particles with magnetic moments being parallel or antiparallel to $\mathbf{B}$, acquire the energy

$$H_f = -\mu B \sum_{i=1}^N s_i. \quad (2)$$

The complete Hamiltonian consists of two terms

$$H = -\frac{J}{N} \sum_{1 \leq i < j \leq N} s_i s_j - \mu B \sum_{i=1}^N s_i. \quad (3)$$

The Hamiltonian (3) does not change if we reverse signs of all spins $s_i \rightarrow -s_i$ ($i = 1, 2, \dots, N$) and the direction of the induction vector $\mathbf{B} \rightarrow -\mathbf{B}$

$$H(s_1, \dots, s_n; \mathbf{B}) = H(-s_1, \dots, -s_n; -\mathbf{B}). \quad (4)$$

Denote a particular configuration $(s_1, s_2, \dots, s_N)$ by $\{s\}$. To each configuration $\{s\}$ there corresponds an energy $E(\{s\}) = H(\{s\})$.

The nature of phase transitions of magnetic systems is well understood. At temperature 0K magnetic systems, in particular the M-W magnet, are in a lowest energy state with all

spins being parallel. Thus, their magnetization M is finite and our magnet is ferromagnetic. As temperature is increased from zero the thermal noise randomizes spins. A fraction of them become antiparallel. This disorder grows with raising temperature and a diminishing fraction of them points at the initial direction. At temperature T_c – the critical temperature, and beyond, magnetization vanishes and the material becomes paramagnetic. For T above T_c , there must be macroscopically large regions in which a net fraction of spins are aligned up. However, their magnetization mutually compensates – they cannot make a finite fraction of all regions agree. For T just below T_c the compensation is not complete and the small, but finite, fraction points in the same direction.

For left-hand vicinity of T_c thermodynamic functions depend on the dimensionless parameter $t = (T - T_c) / T_c$ and consists of terms regular in t and singular in it. The singular terms depend on powers of $|t|^{-1}$. These powers are called the critical indexes (or critical exponents) and are defined for $B = 0$ and $t \rightarrow 0$. The specific heat c per one particle and the magnetic susceptibility χ_T for the paramagnetic phase are characterized respectively by critical indexes α and γ , whereas in the ferromagnetic phase they are characterized by α' and γ' . For example in the paramagnetic phase

$$c \sim |t|^{-\alpha}, \quad \chi_T \sim |t|^{-\gamma}. \quad (5)$$

We shall calculate these sets of critical indexes, as well as the dependence of the internal energy and entropy on $|t|$, for ferromagnetic and paramagnetic phases of the Curie-Weiss magnet.

In the presence of the magnetic field the dependence of magnetization on the magnetic field at $|t| = 0$ is characterized by the critical index δ

$$m \sim B^{1/\delta}. \quad (6)$$

In the following, in place of temperature we will use $\theta = k_B T$, k_B being the Boltzmann constant.

III. CALCULATION OF FREE ENERGY

Note that due to the particular structure of the Hamiltonian of C-W magnet, the Hamiltonian (1) can be written as

$$H_{int} = -\frac{J}{2N} \left(\sum_{i=1}^N s_i \right)^2 + \frac{J}{2}. \quad (7)$$

This form of Hamiltonian can be considered as a defining feature of mean field models.

The partition function is defined as usual as¹¹

$$Z_N = \sum_{\{s\}} e^{-E(\{s\})/\theta}. \quad (8)$$

The summation is performed over all 2^N configurations $\{s\}$. Note that a different method calculating spin configurations can be used³⁻⁵.

Introduce two dimensionless quantities $K = J/\theta$ and $h = \mu B/\theta$. Then, the partition function (8) can be written as

$$\begin{aligned} Z_N(\theta) &= \sum_{\{s\}} \exp \left[\frac{K}{2N} \left(\sum_{i=1}^N s_i \right)^2 - \frac{K}{2} + h \sum_{i=1}^N s_i \right] = \\ &= e^{-\frac{K}{2}} \sum_{\{s\}} \exp \left[\left(\sqrt{\frac{K}{2N}} \sum_{i=1}^N s_i \right)^2 + h \sum_{i=1}^N s_i \right]. \end{aligned} \quad (9)$$

Thermodynamic functions can be obtained via the Helmholtz free energy $F_N(\theta, B)$ ¹¹

$$F_N(\theta, B) = -\theta \ln Z_N(\theta, B). \quad (10)$$

Free energy is minimized when a system reaches equilibrium^{10,11}. The derivative of $F_N(\theta, B)$ with respect to B gives

$$\frac{\partial F_N(\theta, B)}{\partial B} = -\mu \left\langle \sum_{i=1}^N s_i \right\rangle = -\mu \sum_{i=1}^N \langle s_i \rangle, \quad (11)$$

where $\langle s_i \rangle$ means the mean value of s_i calculated with the canonical distribution function¹¹.

The minus derivative (11) defines magnetization M of C-W magnet, thus

$$M = \mu \sum_{i=1}^N \langle s_i \rangle. \quad (12)$$

The second derivative of $F_N(\theta, B)$ with respect to B is positive¹²

$$\frac{\partial^2 F_N(\theta, B)}{\partial B^2} = \frac{\mu^2}{\theta} \left\{ \left\langle \left(\sum_{i=1}^N s_i \right) \left(\sum_{j=1}^N s_j \right) \right\rangle - \left\langle \sum_{i=1}^N s_i \right\rangle^2 \right\} > 0. \quad (13)$$

Since $[\partial^2 F(\theta, B)/\partial B^2]_\theta$ defines the magnetic susceptibility χ_T (cf. Eq. (34)), we conclude that the magnetic susceptibility is positive. Therefore, the change of induction δB increases the magnet energy by $\delta B \cdot \chi \cdot \delta B > 0$. The CW magnet fulfills the condition of thermodynamic (macroscopic) condition of stability¹⁰.

A simple evaluation of the partition function (9) is precluded only by square of magnetization in the exponential. One can get rid of this square using the Gaussian linearization of the form

$$e^{-a^2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} d\xi e^{-\xi^2/2 + \sqrt{2}a\xi}. \quad (14)$$

In the present case $a = \sqrt{K/(2N)} \sum_{i=1}^N s_i$. Now, the partition function factors with respect to individual summations over the state s_i

$$\begin{aligned} Z_N &= \frac{e^{-\frac{K}{2}}}{\sqrt{2\pi}} \sum_{s_1=-1}^{+1} \sum_{s_2=-1}^{+1} \dots \sum_{s_N=-1}^{+1} \\ &\int_{-\infty}^{+\infty} d\xi e^{-\xi^2/2} e^{\sqrt{K/N} s_1 \xi + h s_1} e^{\sqrt{K/N} s_2 \xi + h s_2} \dots e^{\sqrt{K/N} s_N \xi + h s_N} = \\ &= \frac{2^N e^{-K}}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} d\xi e^{-\xi^2/2} \left[\cosh \left(\sqrt{K/N} \xi + h \right) \right]^N. \end{aligned} \quad (15)$$

Performing the change of variable $\sqrt{K/N} \xi = Ky$ we get

$$Z_N = 2^N \left(\frac{KN}{2\pi} \right)^{1/2} e^{-\frac{K}{2}} \int_{-\infty}^{+\infty} dy \Phi_{K,y}(y), \quad (16)$$

where

$$\Phi_{K,h}(y) = e^{-Ky^2/2} \cosh(Ky + h). \quad (17)$$

In addition to y the function $\Phi_{K,h}(y)$ depends on two dimensionless parameters K and h , i.e. on θ and B .

Free energy per one particle is proportional to $(\ln Z_N)/N$. Since we are interested in analyzing the system in the large size limit

$$\lim_{N \rightarrow \infty} \frac{\ln Z_N}{N} = \lim_{N \rightarrow \infty} \ln \left(Z_N^{1/N} \right) = \ln \left(\lim_{N \rightarrow \infty} Z_N^{1/N} \right).$$

For Z_N (15) we obtain

$$-\frac{f(\theta, B)}{\theta} = \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(e^{-\frac{K}{2}} \sqrt{KN/2\pi} \right) + \ln 2 + \ln \lim_{N \rightarrow \infty} \left\{ \int_{-\infty}^{+\infty} dy [\Phi_{K,h}(y)]^N \right\}^{1/N}. \quad (18)$$

In order to obtain the explicit form of the function $f(\theta, B)$ we use the Laplace theorem¹³.

The result is

$$-\frac{f(\theta, B)}{\theta} = \ln \max_{-\infty \leq y \leq \infty} \Phi_{K,h}(y) + \ln 2. \quad (19)$$

Let us introduce a function of y related to free energy

$$f_{\theta,h}(y) = -\theta [\ln 2 + \ln \Phi_{\theta,h}(y)]. \quad (20)$$

To find the dependence of free energy on thermodynamic variables θ and B one should find extreme points of $f_{K,h}(y)$. For these points $(df_{K,h}(y)/dy)_{\theta,B} = 0$. Since

$$\left(\frac{\partial f}{\partial y} \right)_{\theta,B} = -\frac{\theta K}{\Phi_{K,h}(y)} [\tanh(Ky + h) - y] \Phi_{K,h}(y) = -\theta_c [\tanh(Ky + h) - y],$$

the state of minimum of free energy occurs for y obeying the equation

$$y = \tanh(Ky + h). \quad (21)$$

For various values of θ (K) and B (h) solutions of this equation provide the function $y = y(\theta, B)$ of state variables. Therefore, the function $\Phi(\theta, B) = \max_{-\infty \leq y \leq \infty} \Phi_{K,h}(y)$ is a composite function of θ and B , namely $\Phi_{K,y}(y(\theta, B))$ and also depends implicitly on these two state variables. Now Eq. (19) can be rewritten in the form

$$f(\theta, B) = -\theta \ln 2 - \theta \ln \Phi(\theta, B),$$

or

$$f(\theta, B) = -\theta \ln 2 - \theta \ln \left\{ e^{-Ky^2(\theta,B)/2} \cosh[Ky(\theta, B) + h] \right\}. \quad (22)$$

Using two familiar identities¹⁵ $d \tanh x / dx = \cosh^{-2} x$ and

$$\cosh^2 x = (1 - \tanh^2 x)^{-1} = (1 - y^2)^{-1}, \quad (23)$$

we calculate the second partial derivative of f with respect to y

$$\left(\frac{\partial^2 f}{\partial y^2} \right)_{\theta,h} \bigg|_{y=y(\theta,B)} = -\theta K \{ K [1 - y^2(\theta, B)] - 1 \}. \quad (24)$$

In the Appendix we show that there exist solutions of Eq. (21) for which this derivative is positive, hence for them free energy is minimal.

For the reversed magnetic field the solution of Eq. (21) is $-y$

$$-y = \tanh [K (-y - h)]. \quad (25)$$

Therefore, free energy $f(\theta, B)$ (19) is an even function of B (and h)

$$-\frac{f(\theta, -B)}{\theta} = \ln 2 + \ln \left\{ e^{-K(-y(\theta, B))^2/2} \cosh [-K(-y(\theta, B)) - h] \right\} = -\frac{f(\theta, B)}{\theta}. \quad (26)$$

These properties of free energy and solutions of Eq. (21) are the reason why in the following we will always have in mind the positive value of B .

The product of $(\partial y / \partial h)_\theta$ and $(\partial^2 f_{K,h}(y) / \partial y^2)_{\theta,h}$ is positive because

$$\left(\frac{\partial y}{\partial h} \right)_\theta \left(\frac{\partial^2 f_{K,h}(y)}{\partial y^2} \right)_{\theta,h} = \frac{\theta K}{\cosh^2(Ky + h)}. \quad (27)$$

Introduce here $K_c = 1$ – the critical value of the parameter K . From the definition of K it is seen that the critical value of θ is $\theta_c = J$.

In the mean field approximation (cf. for example¹⁴) the Hamiltonian (3) is approximated by

$$H_{mf} = - \sum_{i=1}^N (Jy + \mu B) s_i.$$

For this Hamiltonian the statistical sum is

$$Z_{mf}(\theta, B) = [2 \cosh(Ky + h)]^N$$

and the corresponding free energy is

$$F_{mf}(\theta, B) = -\theta N [\ln 2 + \ln \cosh(Ky + h)].$$

Using the thermodynamic identity (11) we obtain Eq. (21).

IV. FREE ENERGY OF C-W MAGNET IN THE ABSENCE OF THE MAGNETIC FIELD

Suppose that $B = 0$ ($h = 0$). If we plot $g(y) = y$ and $q(y) = \tanh(Ky)$ as functions of y , the points of intersection determine solutions of Eq. (21). Referring to Fig. 1 (left panel),

we have to make distinction between cases. If $\theta > \theta_c$ ($\theta > J$) the slope of the function $q(y)$ at the origin $K = J/\theta = \theta_c/\theta < 1$ is smaller than the slope of linear function $g(y) = y$, which is 1, thus these graphs intersect only at the origin. It is easy to check that for this solution the second derivative of free energy (19) is positive (cf. Appendix). Therefore, the extreme indeed is a minimum (cf. Fig. 2).

On the other hand when $\theta < \theta_c$ ($\theta < J$), the initial slope of $\tanh(Ky)$ is larger than that of linear function, but since values of $\tanh$ function cannot take values outside the interval $(-1, +1)$, the two functions have to intersect in two additional, symmetric nonzero points $\pm y(\theta)$ (Fig. 1). In this case in the Appendix we show that the second derivative of free energy is negative at the origin $y = 0$, which means that there is a maximum at $y = 0$ (cf. Fig. 2). This derivative is positive at $y = \pm y(\theta)$. Free energy $f(\theta)$ attains minimal value at $y = \pm y(\theta)$, hence these solutions correspond to the thermodynamically stable states.




FIG. 1. Graphical solution of Eq. (21). The full line represents the function $g(y) = y$. The dotted lines: $\theta < \theta_c$, the dashed lines: $\theta \geq \theta_c$. Left panel: $h = 0$ ($B = 0$). Right panel: $h > 0$ ($B > 0$).

The parameter K and temperature θ can be expressed in terms of $t = (\theta/\theta_c - 1)$, namely

$$K = \frac{\theta_c}{\theta} = \frac{1}{1+t}, \quad (28)$$

$$\theta = (1+t)\theta_c.$$

V. MAGNETIZATION AND MAGNETIC SUSCEPTIBILITY OF CURIE-WEISS MAGNET

Consider the C-W magnet when the magnetic field B is brought back. Magnetization per particle, m , is a partial derivative of free energy $f(\theta, B)$ after B

$$\begin{aligned} m(\theta, h) &= - \left[\frac{\partial f(\theta, h)}{\partial B} \right]_{\theta} = \mu \left[\frac{\partial \ln \Phi(\theta, h)}{\partial h} \right]_{\theta} = \\ &= \mu \left. \frac{\partial \ln \Phi_{K,h}(y)}{\partial h} \right|_{y=y(\theta, B)} + \mu \left. \frac{\partial \ln \Phi_{K,h}(y)}{\partial y} \right|_{y=y(\theta, B)} \frac{\partial y(\theta, B)}{\partial h}. \end{aligned} \quad (29)$$

Since $\Phi_{K,h}(y)$ attains an extremum at $y(\theta, B)$, the second term on right hand side of (29) vanishes. The contribution of the first term yields the equation of state

$$m = \mu \tanh \left(\frac{K}{\mu} m + h \right), \quad (30)$$

which has a closed analytic form. Therefore, according to Eq. (21), the study of y is equivalent to the study of magnetization m .

Consider the solutions of Eq. (21) for $B > 0$. When $\theta < \theta_c$ the plots of functions $g(y) = y$ and $Q(y) = \tanh(Ky + h)$ intersect in three non symmetric and nonzero points (cf. right panel of Fig. 1). Only for positive value of $y = y(\theta, B)$ free energy f attains the global minimum. To one of negative values there corresponds a local minimum, to the remaining a maximum (cf. Fig. 2). The negative values of $y(\theta, B)$ (as well as $m(\theta, B)$) do not correspond to stable states and should be omitted. One should notice that negative values of y for positive h are not compatible with the symmetry (25) of the state equation. When $\theta \geq \theta_c$ the graphs of $g(y)$ and $Q(y)$ intersect at one point $y(\theta, B) > 0$. For this value of $y(\theta, B)$ free energy has global minimum (cf. Fig. 2).

Consider Eq. (21) for small values of $|t|$ and $h = 0$. We can expand $\tanh(Ky)$ into Taylor's series. Using Eq. (28) we obtain

$$y \simeq \sqrt{3} \frac{(K-1)^{1/2}}{K^{3/2}} \sim |t|^{1/2}. \quad (31)$$

Hence, in agreement with Fig. 3, we conclude that

$$m \sim |t|^{1/2}. \quad (32)$$

When positive $B \rightarrow 0$ then $y(\theta, h) \rightarrow y(\theta)$ of Fig 3. Thus, we note the existence of




FIG. 2. Dependence of function $f_{K,h}(y)$ (19) on y for paramagnetic ($\theta \geq \theta_c$) and ferromagnetic ($\theta < \theta_c$) phases in the magnetic field ($h > 0$) and when the magnetic field is absent ($h = 0$).

spontaneous magnetization $m(\theta) = \mu y(\theta)$

$$m(\theta, h = 0^+) = \begin{cases} 0 & \theta_c \leq \theta, \\ m_0 & \theta_c > \theta. \end{cases} \quad (33)$$

Even after turning off the magnetic field, below critical temperature the system remains magnetized, depending on the sign of B before its removal. The dependence of y for $B = 0$ on θ is singular. It is seen in Fig. 3 that at the point $\theta = \theta_c$ tangents of two branches of the curve are different. We note that the point $\theta = \theta_c$ is the boundary between the region of existence and nonexistence of magnetization, i.e. it is a critical point. We conclude that magnetization m is the order parameter.

In the presence of even an arbitrarily weak ($0 \leq h \ll 1$), the magnetic field magnetization m does not vanish below and above critical point (cf. Fig. 3 the right panel). The external magnetic field lowers the symmetry of the paramagnetic phase. From the point of view of magnetization the difference between paramagnetic and ferromagnetic phases vanishes and

the critical point ceases to exist.




FIG. 3. Dependence of y on temperature. Left panel: $h = 0$. Right panel: $h > 0$.

Consider the magnetic susceptibility

$$\chi_T(\theta, B) = \left[\frac{\partial m(\theta, B)}{\partial B} \right]_{\theta}. \quad (34)$$

According to the definition (34) and Eq. (29) the susceptibility is related to the second derivative of free energy with respect to induction

$$\chi_T(\theta, B) = - \left[\frac{\partial^2 f(\theta, B)}{\partial B^2} \right]_{\theta}. \quad (35)$$

Differentiating both sides of Eq. (30) with respect to B and solving the obtained equation, we obtain the general expression for the susceptibility

$$\chi_T(\theta, B) = \frac{\mu^2}{\theta} \frac{[1 - y^2(\theta, B)]}{1 - K[1 - y^2(\theta, B)]}. \quad (36)$$

Using Eq. (24) the susceptibility can be written as

$$\chi_T(\theta, B) = \mu^2 K \frac{1 - y^2(\theta, B)}{\left(\frac{\partial^2 f_{K,h}(y)}{\partial y^2} \right)_{\theta,h} \Big|_{y=y(\theta,B)}}. \quad (37)$$

Since for a stable state the second derivative $(\partial^2 f_{K,h}(y) / \partial y^2)_{\theta,h} \Big|_{y=y(\theta,B)}$ is nonnegative and $0 \leq |y| \leq 1$, the susceptibility is nonnegative too. Notice that we succeed in linking together the macroscopic (13) and microscopic (36) stability conditions.

For the paramagnetic phase ($\theta \geq \theta_c$ ($t \geq 0$)) $y = 0$, and in the vicinity of the critical temperature we obtain

$$\chi_T = \frac{\mu^2}{\theta_c(1+t)} \frac{(1+t)}{t} = \frac{\mu^2}{\theta_c} t^{-1}. \quad (38)$$

This means that the critical index $\gamma = 1$.

The function $\text{arctanh}y$ obeys the equation

$$\text{arctanh}y = Ky + h. \quad (39)$$

We shall study solutions of Eq. (39) in vicinity of the critical temperature. For small y one can expand $\text{arctanh}y$ into Taylor's series¹⁵. For $\theta < \theta_c$ ($t < 0$) Eq. (39) reduces to

$$y^3 - 3\frac{|t|}{1-|t|}y - 3h = 0. \quad (40)$$

In the ferromagnetic phase and in the absence of the magnetic field, the order parameter does not vanish

$$y^2 = \frac{3|t|}{1-|t|} \quad (t < 0). \quad (41)$$

When $B = 0$ from Eq. (41) it follows that for ferromagnetic phase in vicinity of critical temperature $m \sim \sqrt{3|t|}$. Thus the critical index of magnetization is $\beta' = 1/2$.

Consider the susceptibility (36) in ferromagnetic phase in the vicinity of the critical temperature. Using the expressions (28) and (41) for the ferromagnetic phase we obtain

$$\lim_{B \rightarrow 0} \chi_T(\theta, B) \sim \frac{\mu^2}{\theta_c} |t|^{-1}, \quad (42)$$

and according to the definition (5) for both phases the critical indexes of magnetization are equal: $\gamma = \gamma' = 1$. We shall note that in the case of ferromagnetic phase one should use the relation (40). We conclude that for both phases the magnetic susceptibility of Curie-Weiss magnet is divergent at the critical temperature. This singular behavior is shown in Fig. 4.




FIG. 4. Dependence of the susceptibility on temperature. Left panel: $h = 0$. Right panel $h > 0$.

At the critical point $t = 0$, hence $y = \sqrt[3]{3} h^{1/3}$. Therefore, according the definition (6), the critical index for the critical isotherm $\delta = 3$.

Continuous phase transitions occur when a new state of reduced symmetry develops continuously from the disordered (high temperature) phase. The ordered phase of M-C magnet has lower symmetry than the symmetry (4) of the Hamiltonian, thus the symmetry is spontaneously broken. There exist two equivalent symmetry related states of C-W magnet with magnetization $+m$ i $-m$ respectively, with equal free energies. These states are macroscopically different, so thermal fluctuations will not bring them into contact in the thermodynamic limit. To describe the ordered state we introduced magnetization – the macroscopic order parameter that describes the character and strength of the broken symmetry.

VI. APPROXIMATE THEORY – THE ANALYSIS OF ROOTS OF THE CUBIC EQUATION FOR MAGNETIZATION

Now we shall study the roots of the cubic equation (40) for the ferromagnetic phase. Since free energy is an even function of B we assume that $h > 0$. We shall study the so called the incomplete cubic equation

$$y^3 + 3(-p)y + 2q = 0, \quad (43)$$

where $p = |t| / (1 - |t|)$, $q = -3h/2$.

It is worthwhile to recall that assumptions proposed by Landau for an incompressible magnet result in free energy depending only on even powers of magnetization^{10,16,17}

$$f(m, T) = f_0(T) + \alpha(T) m^2 + \frac{1}{2} \beta(T) m^4.$$

This form of free energy leads to a cubic equation for magnetization.

Introduce a characteristic value of the parameter h_t

$$h_t = \frac{2}{3} \left(\frac{|t|}{1 - |t|} \right)^{\frac{3}{2}}. \quad (44)$$

Roots of Eq. (43) depend on the sign of the discriminant¹⁵ $D = (q^2 - p^3)$

$$D = \left(\frac{3}{2} h \right)^2 - \left(\frac{|t|}{1 - |t|} \right)^3 = \left(\frac{3}{2} \right)^2 (h^2 - h_t^2). \quad (45)$$

If $D < 0$ inequalities $-h_t < h < h_t$ hold. If $D > 0$, then $h > h_t$ or $h < -h_t$.

If $D < 0$ all tree roots are real

$$\begin{aligned} y_1^{(<)}(t, h) &= u_{<}(t, h) + v_{<}(t, h), \\ y_2^{(<)}(t, h) &= \varepsilon_2 u_{<}(t, h) + \varepsilon_1 v_{<}(t, h), \\ y_3^{(<)}(t, h) &= \varepsilon_1 u_{<}(t, h) + \varepsilon_2 v_{<}(t, h), \end{aligned} \quad (46)$$

where

$$\begin{aligned} u_{<}(t, h) &= \sqrt[3]{-q + i\sqrt{|D|}}, \quad v_{<}(t, h) = [u_{<}(t, h)]^*, \\ \varepsilon_1 &= (-1 + i\sqrt{3})/2, \quad \varepsilon_2 = \varepsilon_1^*. \end{aligned} \quad (47)$$

Using relations (47) we can show that

$$\begin{aligned} y_1^{(<)}(t, h) &= 2 \operatorname{Re} u_{<}(t, h), \\ (y_\sigma^{(<)})^*(t, h) &= y_\sigma^{(<)}(t, h) \quad (\sigma = 2, 3). \end{aligned}$$

If $D > 0$ (i.e. $h^2 > h_t^2$) only one root is real

$$y_{>}(t, h) = u_{>}(t, h) + v_{>}(t, h), \quad (48)$$

where

$$u_{>}(t, h) = \sqrt[3]{\frac{3h}{2} + \sqrt{D}}, \quad v_{>}(t, h) = \sqrt[3]{\frac{3h}{2} - \sqrt{D}}. \quad (49)$$

The two remaining two roots are complex.

If we combine Eqs. (44)-(49) we obtain the functions $u_{<}(t, u)$, $v_{<}(t, u)$ and $u_{>}(t, u)$, $v_{>}(t, u)$ in the useful form

$$\begin{aligned} u_{>}(t, h) &= (3/2)^{1/3} \sqrt[3]{h + \sqrt{h^2 - h_t^2}}, \quad v_{>}(t, h) = (3/2)^{1/3} \sqrt[3]{h - \sqrt{h^2 - h_t^2}}, \\ u_{<}(t, h) &= (3/2)^{1/3} \sqrt[3]{h + i\sqrt{h_t^2 - h^2}}, \quad v_{<}(t, h) = (3/2)^{1/3} \sqrt[3]{h - i\sqrt{h_t^2 - h^2}}. \end{aligned} \quad (50)$$

Consider functions $u_{>}(t, h)$ and $v_{>}(t, h)$ for negative h

$$\begin{aligned} u_{>}(t, -h) &= \sqrt[3]{-1} \sqrt[3]{h - \sqrt{h^2 - h_t^2}} = \sqrt[3]{-1} v_{>}(t, h), \\ v_{>}(t, -h) &= \sqrt[3]{-1} \sqrt[3]{h + \sqrt{h^2 - h_t^2}} = \sqrt[3]{-1} u_{>}(t, h). \end{aligned} \quad (51)$$

Similar relations hold also for $u_{<}$ and $v_{<}$. Among three root of unity ε_1 , ε_2 and $\varepsilon_3 = -1$ only the latter root yields proper symmetry relation (25)

$$y_j^{(\sigma)}(t, -h) = -y_j^{(\sigma)}(t, h) \quad (\sigma = >, <; j = 1, 2, 3). \quad (52)$$

In Fig. 5 we have plotted the dependence of roots (46) and (49) on the magnetic field h . For positive values of $y_{<}^{(1)}$ and $-h_t < h < 0$, as well as for negative values of $y_{<}^{(3)}$ and $0 < h < h_t$, the signs of h and these two roots do not agree. The symmetry (25) of equation of state is broken. On the line BB' (corresponding to the root $y_{<}^{(2)}$) the derivative $(\partial y(\theta, h)/\partial h)_\theta < 0$ is negative. From Eq. (27) it follows that $(\partial^2 f_{\theta, h}(y)/\partial y^2)_{\theta, h}$ is negative too. In this interval of values of h free energy has maxima, therefore CW magnet is not in stable states. We conclude that for the root $y_{<}^{(2)}(t, h)$ free energy attains maximal values.




FIG. 5. Plot of dependence of roots of Eq. (43) on h for $h_t = 0.1$. For $|h| > h_t$ the curve represents the function $y_{>}(t, h)$. In the interval $-h_t < h < h_t$ the plot consists of $y_{<}^{(m)}(t, h)$ ($m = 1, 2, 3$)

For $0 < h < h_t$ the root $y_{<}^{(1)}$ is positive. The remaining two roots are negative and $|y_{<}^{(3)}| < y_{<}^{(1)}$ (cf. Fig. 6). Further $y_{<}^{(2)} \rightarrow y_{<}^{(3)}$ when $h \rightarrow h_t^- \equiv h_t - \varepsilon$; ($0 < \varepsilon \ll 1$) and

$$\lim_{h \rightarrow h_t^-} \left(\frac{\partial y_{<}^{(2)}(\theta, h)}{\partial y} \right)_\theta = -\infty.$$




FIG. 6. Dependence of $|y_{<}^{(3)}|$ and $y_{<}^{(1)}$ on h .

From Eq. (27) it follows that

$$\lim_{h \rightarrow h_t^-} \left(\frac{\partial^2 f_{\theta,h}(y)}{\partial y^2} \right)_{\theta,h} = 0.$$

This means that for $h = \pm h_t$ one should consider the third derivative of free energy with respect to y .

In the interval $-h_t < h < h_t$ the derivative of the root $y_{<}^{(3)}$ is positive, therefore the derivative $(\partial^2 f_{\theta,h}(y) / \partial y^2)_{\theta,h} \big|_{y=y_{<}^{(3)}}$ is also positive. This means that in this interval free energy has a (local) minimum. Free energy exhibits the global minimum for the root $y_{<}^{(1)}$ for $h_t > h > 0$. This behavior of free energy and of solutions of cubic equation is shown in Fig. 5.

For $h > h_t$ and $h < -h_t$ free energy has the global maximum for the real root $y_{>}(t, h)$ (cf. Fig. 5).

Note that as a result of approximation yielding Eq. (40) values of $|y|$ may exceed the limiting value 1. This means that we shall restrict ourselves to small values of $|t|$ and h_t .

We shall point out that value h_t (44) of parametr h define such value B_t of magnetic induction B , for which the value of induced magnetization $m_{ind} \sim \chi_T \cdot B_t$ ($B_t = \theta_c h_t / \mu$ with h_t given by Eq. (38)), is in accordance with Eq. (41).

If $h \ll h_t$ ($|t| \neq 0$) the magnetic field B is weak and does not influence the thermodynamic quantities characterizing the system. If $h \gg h_t$ the field B is strong. If $t = 0$ ($T = T_c$) all magnetic fields are strong. As we have shown, if $t \sim 0$ and field is strong, $m \sim h^{1/3}$.

VII. PROPERTIES OF THE INTERNAL ENERGY, ENTROPY AND SPECIFIC HEAT OF THE CURIE-WEISS MAGNET

To find the internal energy U we shall use the familiar thermodynamic identity¹¹

$$U = -\theta^2 \frac{\partial}{\partial \theta} \left(\frac{F}{\theta} \right) = N\theta^2 \frac{\partial}{\partial \theta} \left(-\frac{f}{\theta} \right)_B.$$

For the internal energy per one spin this formula gives

$$u = -\theta^2 \frac{\partial}{\partial \theta} \left[-Ky^2/2 + \ln \cosh (Ky + h) \right].$$

As a result of simple calculations we obtain

$$u(\theta, B) = -\frac{J}{2}y^2(\theta, B) - \mu By. \quad (53)$$

The first term of this equation is the interaction energy per one spin, whereas the second term is the energy of a spin in the magnetic field. When $\theta = 0$, $y = 1$, hence $u = -J/2 - B\mu$. When $B = 0$, in the paramagnetic phase $y = 0$ and $u = 0$. When $B = 0$ in the ferromagnetic phase in vicinity of θ_c one has $K \sim (1 - |t|)^{-1}$ and

$$y \approx \pm \sqrt{3|t|/(1 - |t|)}, \quad (54)$$

thus, $u \sim |t|^1$. As we can see from Fig. 7, in the absence of the magnetic field the behavior of internal energy is singular at $K = 1$. The curve representing the function $u(\theta, h = 0)$ consist of two branches. At $\theta = \theta_c$ their derivatives are different. When the magnetic field is turned in this singularity is washed out.

Consider entropy

$$S = -Nk_B \left(\frac{\partial f(\theta, B)}{\partial \theta} \right)_B. \quad (55)$$

Calculating the derivative we obtain for entropy per particle the familiar thermodynamic identity¹¹

$$s(\theta, B) = \frac{k_B}{\theta} [-f(\theta, B) + u(\theta, B)]. \quad (56)$$




FIG. 7. Dependence of the internal energy on temperature. Left panel: $h = 0$. Right panel $h > 0$.

For low temperatures

$$f(\theta, B) \sim \frac{u(\theta, B)}{\theta}.$$

Therefore, even when the magnetic is turned in, at $\theta = 0$ entropy vanishes. The spins are completely ordered, therefore entropy acquires the lowest value. Since in the absence of the magnetic field the internal energy is singular at K_c , entropy is also singular at the critical point. When $B = 0$ in the paramagnetic phase ($\theta \geq \theta_c$) $y = 0$, and the internal energy vanishes. From the definition (19) it follows that $-k_B f(\theta \geq \theta_c, B = 0)/\theta = k_B \ln 2$, and $s = k_B \ln 2 \simeq 0.7 \times k_B$. Spins in the paramagnetic phase are completely disordered and entropy reaches its greatest value. The dependence of entropy on temperature is shown in Fig. 8.




FIG. 8. Dependence of entropy on temperature. Left panel: $h = 0$. Right panel: $h > 0$.

If $B = 0$ in the paramagnetic phase $y = 0$, hence $u(\theta > \theta_c, h = 0) = 0$. In the ferromagnetic phase and in vicinity of θ_c according (31) $u \sim |t|^1$. For $\theta = \theta_c$ the internal energy vanishes $u = 0$. This means that the internal energy is the continuous function of

temperature $u(\theta_c^-) = u(\theta_c^+) = 0$.

Entropy s also is a continuous function of temperature. To show this property in the case of ferromagnetic phase and vicinity of θ_c we use Eqs. (22) and (31) and Tylor's series for $\ln [\tanh (Ky)]$ we get

$$s \approx k_B (\ln 2 - 3 |t| / 2) .$$

We see that, as one may expect, that entropy of the ferromagnetic phase is smaller than entropy of paramagnetic phase and for $|t| = 0$ attains its maximal value.

The behavior of heat capacity in vicinity of the critical temperature is more complex. To calculate heat capacity per one particle one can use one of two thermodynamic relations, namely¹¹

$$c(\theta, B) = k_B \left(\frac{\partial u(\theta, B)}{\partial \theta} \right)_B , \quad (57)$$

or

$$c(\theta, B) = \theta \left(\frac{\partial s(\theta, B)}{\partial \theta} \right)_B . \quad (58)$$

It is an easy task to show that these identities yield the same result. We shall consider Eq. (57). From Eq (53) it follows that heat capacity $c(\theta, B)$ depends on derivative $(\partial y / \partial \theta)_B$

$$c = -k_B \theta (Ky + h) \left(\frac{\partial y}{\partial \theta} \right)_B .$$

Differentiating both sides of Eq. (21) with respect to y , solving the obtained equation for $(\partial y(\theta, B) / \partial \theta)_B$ and applying the identity (23), we obtain an analytic expression for this derivative

$$\left(\frac{\partial y}{\partial \theta} \right)_B = -\frac{1}{\theta} \frac{(Ky + h)}{\cosh^2(Ky + h) - K} .$$

With the help of the above relation we obtain the final form of expression for heat capacity per one particle

$$c = k_B \frac{(1 - y^2) (Ky + h)^2}{(1 - K) + Ky^2} . \quad (59)$$

When $B = 0$ in the paramagnetic phase ($\theta \geq \theta_c$) magnetization m vanishes, i.e. $y = 0$. Hence, $c = 0$. Since $\lim_{\theta \rightarrow 0} y = 1$ in this limit c also vanishes.

Consider heat capacity in ferromagnetic phase in the vicinity of the critical temperature. Using expressions (28) and (54) we obtain

$$\frac{c}{k_B} \approx \frac{3}{2} (1 - |t|) . \quad (60)$$

Heat capacity is discontinuous at θ_c

$$c(\theta_c^-, B=0) - c(\theta_c^+, B=0) = 3k_B/2, \quad (61)$$

with $\theta_c^\pm = \theta_c \pm \epsilon$, and $0 < \epsilon \ll 1$. The zero field heat capacity is singular at the critical temperature, whereas in the magnetic field it exhibits a peak at the transition point (cf. Fig. 9).




FIG. 9. Dependence of heat capacity per one particle on temperature. Left panel: $h = 0$. Right panel: $h > 0$.

VIII. NUMERICAL RESULTS

Until now we described the phase transition using the approximate expression for $\text{arctanh}(Ky + h)$. Now we shall give up this approximation, which means that we shall rely on numerical calculations. Such approach will shed light on the approximation used in Sect. VI.

Let us rewrite Eq. (39) in the form

$$\text{arctanh} y - \frac{y}{1 - |t|} = h. \quad (62)$$

For a given value of $|t|$ we numerically solve Eq. (62) for various values of h . We plot $y = y_t(h)$ in Fig. 10. The obtained plot resembles the plot obtained for roots of Eq. (43). However, the critical values of the parameter $h_t^{(num)}$ are smaller than obtained $h_t^{(approx)}$ as a result of calculation based on the approximate theory presented in the Sect. VI. Besides, unlike for the approximate theory, values of y_{num} are confined in the interval $[-1,1]$. In Fig.




FIG. 10. Dependence of $y_t(h)$ on the magnetic field h . Dashed line represents the dependence of roots of Eq. (43) on the magnetic field. The full line is plot of the function (62).

11 we compare values of $h_t^{(approx)}$ and $h_t^{(num)}$. It is seen that for small values of $|t|$ both plots differ a little. With growing $|t|$ difference is more pronounced.

IX. APPENDIX

We shall show that there exist solutions $y(\theta, B)$ of Eq. (39) for which the second derivative of function $f_{K,h}(y)$ (Eq. (20)) with respect to y calculated at $y = y(\theta, B)$

$$\left(\frac{\partial^2 f_{\theta,h}(y)}{\partial y^2} \right)_{\theta,h} \Big|_{y=y(\theta,B)} = -\theta K \{ K [1 - y^2(\theta, B)] - 1 \} \quad (63)$$

is positive.

Consider temperatures higher than θ_c .

1. For $B = 0$ in the paramagnetic phase $K \leq 1$ and magnetization vanishes. In this case $y = 0$, therefore, the second derivative (63) $[-\theta K (K - 1)] > 0$ is positive.




FIG. 11. Plot of dependence of critical value of the magnetic field h_t on the parameter t . The full line – result numerical calculation, the dashed line represents the dependence resulting from the cubic equation (43).

2. If $B > 0$ magnetization is positive, hence $1 > y > 0$. The double inequality $0 < K[1 - y^2(\theta, B)] < 1$ holds. Therefore, $K[1 - y^2(\theta, B)] - 1 < 0$. Hence, the second derivative (63) is positive. The same arguments are valid for the negative value of induction ($B < 0$ and $y < 0$).

In the case of $\theta < \theta_c$ the parameter K is greater than unity. As we know (cf. Sect. IV), if $B = 0$, Eq. (21) has three solutions, namely $y = 0$ and $y = \pm y(\theta) \equiv \pm y_0$.

1. For $y = 0$ the derivative (63) is negative

$$\left(\frac{\partial^2 f_{\theta,h}(y)}{\partial y^2} \right)_{\theta,h} \bigg|_{y=0} = \theta K (1 - K) < 0. \quad (64)$$

This means that in the ferromagnetic phase the solution $y = 0$ corresponds to a maximum of free energy.

2. In the case of two remaining solutions the second derivative reads

$$\left(\frac{\partial^2 f_{K,h}(y)}{\partial y^2} \right)_{\theta,h} \bigg|_{y=\pm y_0} = -\theta K [K(1 - y_0^2) - 1]. \quad (65)$$

We shall express the parameter K by y_0 . Using Eq. (39) we can write

$$K = \frac{\operatorname{arctanh} y_0}{y_0}.$$

With the help of this relation we find

$$\left(\frac{\partial^2 f_{K,h}(y)}{\partial y^2} \right)_{\theta,h} \Big|_{y=\pm y_0} = -\theta K \left[\frac{\operatorname{arctanh} y_0}{y_0} (1 - y_0^2) - 1 \right]. \quad (66)$$

For small y_0 we can use the approximate expression $\operatorname{arctanh} y_0 \approx y_0 + y_0^3/3$. This yields the inequality

$$\frac{(1 - y_0^2) \operatorname{arctanh} y_0}{y_0} \approx (1 + y_0^2/3) (1 - y_0^2) < (1 - y_0^4) < 1,$$

and the second derivative of free energy is positive. In the ferromagnetic phase, in vicinity of critical temperature, for solutions $\pm y_0$ free energy reaches a minimum.

The function

$$\varphi(y_0) = \frac{\operatorname{arctanh} y_0}{y_0} (1 - y_0^2)$$

is monotonically decreasing in the interval $(0, 1]$ and $0 \leq \varphi(y_0) < 1$.

Using the logarithmic representation¹⁸ $\operatorname{arctanh} y_0 = \frac{1}{2} \ln \left(\frac{y_0+1}{y_0-1} \right)$ and the limiting value¹⁸ $\lim_{x \rightarrow 0} x \ln x = 0$ we can show that

$$\lim_{y_0 \rightarrow 1^-} \frac{\operatorname{arctanh} y_0}{y_0} (1 - y_0^2) = 0,$$

hence, for low temperatures the right hand side of Eq. (66) is positive. The plot of function (66) is shown in Fig. 12. We conclude that in the ferromagnetic phase minimums of free energy occur for $y = \pm y_0$.

3. Assume that $B > 0$ ($h > 0$) and $\theta < \theta_c$ ($K > 1$). In this case, according to Sect. V, Eq. (21) has three solutions $y_1^{(<)}(t, h) > 0$, $y_2^{(<)}(t, h) < 0$, $y_3^{(<)}(t, h) < 0$. Proceeding as before we express K by $y_1^{(<)}$ and h

$$K = \frac{\operatorname{arctanh} y_1^{(<)} - h}{y_1^{(<)}}. \quad (67)$$

Now, Eq. (63) takes the form

$$\left(\frac{\partial^2 f_{K,h}(y)}{\partial y^2} \right)_{\theta,B} \Big|_{y=y_1^{(<)}} = -\theta K \left\{ \frac{\operatorname{arctanh} y_1^{(<)} - h}{y_1^{(<)}} \left[1 - \left(y_1^{(<)} \right)^2 \right] - 1 \right\}. \quad (68)$$




FIG. 12. Plot of $f_{K,h}(y)$ (solid line), $df_{K,h}(y)/dy$ (broken line) and $d^2f_{K,h}(y)/dy^2$ (dotted line). To zeros of $df_{K,h}(y)/dy$ there correspond extremums of $f_{K,h}(y)$. Two of them are minimums because $\frac{d^2f_{K,h}(y)}{dy^2} > 0$.

Since $0 < y_1^{(<)} \leq 1$, the inequality $\left[1 - \left(y_1^{(<)}\right)^2\right] < 1$ holds. Thus, the additional term of Eq. (68) is non-positive

$$-\frac{h}{y_1^{(<)}} \left[1 - \left(y_1^{(<)}\right)^2\right] \leq 0,$$

and, even in this case, the second derivative of free energy is positive.

* tapasz@prz.edu.pl

- ¹ S-K Ma, *Modern Theory of Critical Phenomena* (Benjamin, Reading, MA, 1976).
- ² S. Adams, *Lectures on Mathematical Statistical Mechanics*, Dublin Institute for Advanced Studies, 2006.
- ³ A. Bovier and I. Kurkova, *A Short Course on Mean Field spin glasses*.
- ⁴ R. Kühn, *Equilibrium Analysis of Complex Systems, Lecture Notes of 7CCMCS03*, King's College, London 2010.
- ⁵ N. Macris, *Statistical Physics for Communication and Computer Science, Lecture Notes 3: Ising Model*, École Polytechnique Fédérale de Lausanne, 2011.
- ⁶ M. Kac, "Mathematical Mechanisms of Phase Transitions" in: *Statistical Mechanics of Phase Transitions and Superfluidity*, edited by M. Chrétinin, E.P. Gross, S. Dresser (Gordon and Breach Science Publishers, New York, 1968).
- ⁷ N. Merhav, *Statistical Physics and Information Theory*, Foundations and Trends in Communications and Information Theory, **6**, 1212 (2009).
- ⁸ A. Montanari, *Stat 316, Stochastic Processes on Graphs, Introduction and the Curie-Weiss model* Stanford University, 2007.
- ⁹ H. Nishimori, *Statistical Physics of Spin Glasses and Information Processing. An Introduction* (Clarendon Press, Oxford, 2001).
- ¹⁰ L.E. Reichl, *A Modern Course in Statistical Physics*, Chapter 4, (University of Texas Press, Austin, 1980).
- ¹¹ K. Huang, *Statistical Mechanics* (Wiley, Hoboken NJ, 1987).
- ¹² F. Reif, *Statistical Physics, Berkeley Physics Course*, Vol. 5 (McGrawHill, New York, 1967).
- ¹³ G. Pólya, G. Szegő, *Problems and Theorems in Analysis I: Series, Integral Calculus, Theory of Functions*, Chapter 5 (Springer, Heidelberg, 1976).
- ¹⁴ P. Palffy-Muhoray, Am. J. Phys. **70**, 433, 2002.
- ¹⁵ I.N. Bronshtein, K.A. Semendyaev, G. Musiol, H. Muehling, *Handbook of Mathematics* (Springer, 4th edition 2004).
- ¹⁶ J. Als-Nielsen, R.J. Birgenau, Am. J. Phys. **45**, 554, 1977.
- ¹⁷ J.J. Binney, N.J. Dowrick, A.J. Fisher, M.E.J. Newman, *The Theory of Critical Phenomena. An Introduction to Renormalization Group* (Clarendon Press, Oxford 1992).
- ¹⁸ *Handbook of Mathematical functions* edited by Milton Abramowitz and Irene A. Stegun (Dover Publications, New York, 1964).